



PCT/GB 2003 / 0 0 3 0 8 3

Der' CT/PTG

14 JAN 2005

GB03/03083

INVESTOR IN PEOPLE

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

REC'D 13 AUG 2003

WIPO

PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.


Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

Dated

4 August 2003

BEST AVAILABLE COPY



28NOV02 1766961-6 002819
P017700 0.00/0227740.8

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

1. Your reference

C245.00/Q

2. Patent application number

(The Patent Office will fill in this part)

0227740.8

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Quest International B.V.
Huizerstraatweg 28
1411 GP Naarden
The Netherlands

Patents ADP number (if you know it)

4329587002

If the applicant is a corporate body, give the country/state of its incorporation

The Netherlands

4. Title of the invention

Improvements in or relating to
perfume compositions

5. Name of your agent (if you have one)

Keith W Nash & Co

"Address for service" in the United Kingdom
to which all correspondence should be sent
(including the postcode)

90-92 Regent Street
Cambridge
CB2 1DP
United Kingdom

Patents ADP number (if you know it)

1206001

If you are declaring priority from one or more
earlier patent applications, give the country
and the date of filing of the or of each of these
earlier applications and (if you know it) the or
each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

UK

0216633.8

18.07.2002

If this application is divided or otherwise
derived from an earlier UK application,
give the number and the filing date of
the earlier application

Number of earlier application

Date of filing
(day / month / year)

Is a statement of inventorship and of right
to grant of a patent required in support of
this request? (Answer "Yes" if:

Yes

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

| | | |
|-------------|----|-----|
| Description | 26 | / |
| Claim(s) | 3 | / w |
| Abstract | 1 | / |
| Drawing(s) | | |

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

1. I/We request the grant of a patent on the basis of this application.
Signature Keith W Nash & Co. Date 27.11.2002
Keith W Nash & Co., Agents

2. Name and daytime telephone number of person to contact in the United Kingdom
Mrs H C Matthews (01223) 355477

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.

Write your answers in capital letters using black ink or you may type them.

If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.

If you have answered 'Yes' Patents Form 7/77 will need to be filed.

Once you have filled in the form you must remember to sign and date it.

For details of the fee and ways to pay please contact the Patent Office.

C245.00/Q

Title: Improvements in or relating to perfume compositions

Field of the Invention

This invention relates to perfume compositions, to products containing such perfume compositions, and to the use of a perfume component or perfume composition to deliver a deodorant effect. In particular, the invention relates to perfume components, mixtures thereof, and perfume compositions for reducing or preventing body malodour.

Background to the Invention

It is known that, at the point of secretion, sweat is odourless. Body malodour is the result of a variety of biotransformations of components of sweat by certain species of natural micro-organisms which live on the surface of the skin. These transformations produce a number of volatile odoriferous compounds such as steroidal compounds (e.g. 16-androstenes), amongst others, which contribute to body malodour.

There are three types of personal product routinely used to combat body malodour: perfumes, antiperspirants and deodorants.

Perfumes may simply mask body malodour. However perfume compositions have been disclosed which exhibit a deodorant action. EP-B-3172, EP-A-5618, US-A-43044679, US-A-4322308, US-A-4278658, US-A-4134838, US-A-4288341 and US-A-4289641 all describe perfume compositions which exhibit a deodorant action when applied to human skin or when included in a laundry product used to launder textiles.

Antiperspirants work by blocking the sweat glands, thereby reducing perspiration.

Antimicrobial agents used in deodorants are designed to reduce the population, inhibit the

growth or diminish the metabolic activities of micro-organisms living on the surface of the skin. Typical agents of this nature include ethanol and Triclosan (2',4,4'-trichloro-2-hydroxydiphenyl ether) which are well known to exert antimicrobial effects. The use of common deodorant actives results in a non-selective antimicrobial action exerted upon most of the skin's natural microflora. This is an undesirable disadvantage of such deodorant formulations, since the natural microflora provides a protective barrier (colonisation resistance) against invasion by potentially pathogenic bacteria.

US-A-5643559 (Colgate-Palmolive Company) discloses deodorant active materials having an effective amount of Zn^{2+} ions for inhibiting bacterial exoenzymes responsible for the production of axillary malodour. The bacterial exoenzymes are further characterised as aryl sulphatase or beta glucuronidase.

DE-4343265 (Henkel) describes deodorant compositions comprising saturated dioic acid (C3-C10) esters. The active inhibits a sweat decomposing esterase and the compositions are said not to disturb the skin's natural microflora.

WO 94/07837 (Unichema) describes certain novel unsaturated dioic acids having between 8 and 22 carbon atoms. The potential use of these acids to treat malodour is also described.

Gower et al. (*J Steroid Biochem. Molec. Biol.*, (1994) Vol. 48, No. 4, pp 409-418) discloses the importance of certain bacterial enzymes involved in bacterial steroid metabolism in the production of odoriferous steroids, and proposes a series of interconversions between some of these metabolites.

Talalay, P.: Hydroxysteroid Dehydrogenases in *The Enzymes*, VII, 2nd Ed., (Boyer, P., Lardy, H., and Myrback, K., eds.), Academic Press, NY, 177, 1963, describes that 3[[alpha]] hydroxysteroid dehydrogenase from *Pseudomonas testosteroni* is inhibited by heavy metals and sulfhydryl-binding reducing agents.

Nakajin et al. (*J Steroid Biochem. Molec. Biol.*, (1991) Jan;38(1):95-9) discloses that the -conazole antifungal agents have a mode of action based on the inhibition of sterol metabolism. The activity of the enzyme (16-ene-C19-steroid synthesizing enzyme) responsible for the conversion of C21-steroids to 16-ene-C19-steroids, which was localized on pig testicular microsomes, was inhibited by some typical imidazole antifungal compounds such as clotrimazole, econazole, miconazole and ketoconazole which are known to be universal inhibitors of cytochrome P-450 dependent enzymes.

Lavallee et al. (*J Steroid Biochem. Molec. Biol.* (1993) Jul;46(1):73-83) describes 20 beta-hydroxypregnenolone as a more potent inhibitor of 5,16-androstadien-3 beta-ol synthetase than of 17-hydroxylase and for the latter enzyme activity, the $K_i(\text{app})$ was lower than that for 17-hydroxypregnenolone itself.

Watabe et al. (*J Biol. Chem.* (1985) Jul 25;260(15):8716-20) describes that the C16-double bond of the steroid androsta-5,16-dien-3 beta-ol, is oxidized by male rat liver microsomes to 16 alpha,17 alpha-epoxyandrost-5-en-3 beta-ol; 16 beta,17 beta-epoxyandrost-5-en-3 beta-ol; androst-5-ene-3 beta,16 alpha,17 beta-triol; and androst-5-ene-3 beta,16 beta,17 alpha-triol, and this transformation is strongly inhibited with CO.

WO 00/01355 and WO 00/01358 describe agents useful in preventing or reducing body malodour by inhibiting the production of odoriferous steroids, wherein the agents inhibit the bacterial enzymes, bacterial 4-ene reductase and/or 5 α -reductase. Examples of active agents are described as dicarboxylic acids, phenyl compounds, monoterpene derivatives, sterols, flavonoids, steryl esters, 2,7-napthalenediol and oxyquinoline (WO 00/01355), and certain perfume components (WO 00/01358).

Several steroids, notably 5 α -androst-16-en-3-one (5 α -androstenone), 5 α -androst-16-en-3 α -ol (3 α -androstenol) and androsta-4,16-dien-3-one (androstadienone) are known to be highly odorous in the context of human axillary odour. The biotransformations effected by a micro-organism on the components of sweat to produce such odoriferous products or

intermediates, occur via a number of possible, and typically, ill-defined metabolic pathways.

It has been suggested in the prior art (Gower et al) that odorous steroids e.g. androstenones are formed by the biotransformation of typically non-odorous steroids i.e. steroids present at levels below the threshold of human olfactory detection, by the action of micro-organisms present on the skin surface. More particularly, 5 α -androsta-5,16-dien-3 β -ol (androstadienol) was a source of the odorous androstenones.

Summary of the Invention

The present invention is based on extensive testing of perfume components to determine whether a particular component is capable of inhibiting the biotransformation of androstadienols to androstenones, particularly 5 α -androst-16-en-3-one, and thus is capable of inhibiting the production of odoriferous steroids by micro-organisms on the skin surface. Based on this testing, perfume components were identified, which whilst known, possess hitherto unappreciated deodorant properties. The invention thus enables perfume compositions to be defined that reduce or prevent body malodour.

Accordingly, in one aspect, the present invention provides a perfume composition comprising at least 30% by weight of one or more of the following perfume components: Armoise Tunisian oil, para-tert.butylphenylacetonitrile, dihydrolinalol, N-ethyl-N-(3-methylphenyl)propionamide, 4-(5-ethylbicyclo[2.2.1]heptyl-2)-cyclohexanol, ethyltricyclo[5.2.1.0{2,6}]decane-2-carboxylate, 2-ethyl-N-methyl-N-(3-methylphenyl)butanamide, dihydromyrcenol, (4-isopropylcyclohexyl)methanol, 3-methyl-5-phenylpentan-1-ol, 2,2,2-trichloro-1-phenylethyl acetate, isobornyl acetate, allyl amyl glycolate, alpha-terpineol, acetyl cedrene, tetrahydrogeraniol, citronellal, cuminic aldehyde, 1,3,3-trimethyl-2-norbornanyl acetate, cis-jasmone, methyl octyl acetaldehyde, gamma-octalactone, octyl acetate, pine American oil, peppermint (Chinese), 1,3,3-trimethyl-2-norbornanol, gamma-nonolactone, octahydro-2H-chromen-2-one, cis-4-decenal, 3-(3-isopropylphenyl)butanal.

The following perfume components are useful in the perfume compositions defined herein:

Armoise Tunisian oil;

para-tert-Butylphenylacetonitrile (also known as 'Marenil' where MARENIL is a trade mark of Quest International);

Dihydrolinalol (3,7-dimethyloct-6-en-3-ol);

N-Ethyl-N-(3-methylphenyl)propionamide (also known as 'Agarbois' where AGARBOIS is a trade mark of Quest International);

4-(5-Ethylbicyclo[2.2.1]heptyl-2)-cyclohexanol;

Ethyltricyclo[5.2.1.0{2,6}]decane-2-carboxylate;

2-Ethyl-N-methyl-N-(3-methylphenyl)butanamide (also known as 'Paradisamide' where PARADISAMIDE is a trade mark of Quest International) ;

Dihydromyrcenol (2,6-dimethyl-7-octen-2-ol);

(4-isopropylcyclohexyl)methanol;

3-Methyl-5-phenylpentan-1-ol (also known as 'Mefrosol' where MEFROSOL is a trade mark of Quest International);

2,2,2-Trichloro-1-phenylethyl acetate (also known as Rosacetone or Roseacetone);

Isobornyl acetate;

Allyl amyl glycolate ('2-methylbutyloxyacetic acid, 2-propenyl ester');

alpha-Terpineol;

Acetyl cedrene (also known as 'Lixetone' where LIXETONE is a trade mark of Quest International);

Tetrahydrogeraniol;

Citronellal;

Cuminic aldehyde (para-isopropylbenzaldehyde);

cis-Jasmone;

Methyl octyl acetaldehyde (2-methyldecenal);

gamma-octalactone (5-butyldihydrofuran-2(3H)-one);

Octyl acetate;

Pine American oil;

Peppermint (Chinese);

1,3,3-Trimethyl-2-norbornanyl acetate (fenchyl acetate);

1,3,3-Trimethyl-2-norbornanol (fenchyl alcohol);

gamma-Nonalactone;

Octahydro-2H-chromen-2-one (also known as 'Octahydrocoumarin' where OCTAHYDROCOUMARIN is a trade mark of Quest International);

cis-4-Decenal;

3-(3-isopropylphenyl)butanal.

The term "perfume component" is used herein to represent a material which is added to a perfume composition to contribute to the olfactive properties of the composition. A perfume component can be acceptably employed to provide odour contributions to the overall hedonic performance of products. Typically, a perfume component will be generally recognised as possessing odours in its own right, will be relatively volatile and often has a molecular weight within the range 100 to 300. Typical materials which are perfume components are described in "Perfume and Flavour Chemicals", Volumes I and II (Steffan Arctander, 1969).

For the purposes of the present invention, by perfume composition is meant a mixture of individual perfume components, and optionally one or more suitable diluents, which is used to impart a desired odour to the skin and/or product for which an agreeable odour is indispensable or desirable. Commonly used diluents are benzyl benzoate, diethyl phthalate, dipropylene glycol and isopropyl myristate. The concentration of perfume components referred to herein is relative to the total concentration of perfume components present in the composition, i.e. excludes any diluents.

To deliver high deodorant effects the perfume component(s) are preferably present in a perfume composition in an amount of 40% by weight of the total weight of the perfume composition, more preferably at least 45%, and most preferably at least 60%.

Additionally, or alternatively, a perfume composition in accordance with the present invention preferably comprises at least 3, more preferably at least 5, and even more preferably at least 10 of the specified perfume components.

Thus, in a further aspect, the present invention provides a perfume composition comprising at least 3 of the following perfume components; Armoise Tunisian oil, para-tert.butylphenylacetonitrile, dihydrolinalol, N-ethyl-N-(3-methylphenyl)propionamide, 4-(5-ethylbicyclo[2.2.1]heptyl-2)-cyclohexanol, ethyltricyclo[5.2.1.0{2,6}]decane-2-carboxylate, 2-ethyl-N-methyl-N-(3-methylphenyl)butanamide, dihydromyrcenol, (4-isopropylcyclohexyl)methanol; 3-methyl-5-phenylpentan-1-ol, 2,2,2-trichloro-1-phenylethyl acetate; isobornyl acetate; allyl amyl glycolate, alpha-terpineol, acetyl cedrene, tetrahydrogeraniol, citronellal, cuminic aldehyde, 1,3,3-trimethyl-2-norbornanyl acetate, cis-jasmone, methyl octyl acetaldehyde, gamma-octalactone, octyl acetate, pine American oil, peppermint (Chinese), 1,3,3-trimethyl-2-norbornanol, gamma-nonolactone, octahydro-2H-chromen-2-one, cis-4-decenal, 3-(3-isopropylphenyl)butanal.

The perfume components useful herein in a perfume composition may be incorporated into deodorant products which include, but are not limited to, body deodorants and antiperspirants including roll ons, sprays, gel products, stick deodorants, antiperspirants, shampoos, soaps, shower gels, talcum powder, hand creams, skin conditioners, sunscreens, sun tan lotions, skin and hair conditioners.

Thus, in an even further aspect, the present invention provides a deodorant product comprising a perfume composition in accordance with the invention.

A deodorant product preferably comprises at least 0.05% to 4%, more preferably 0.1% to 2% of a perfume composition by weight of the deodorant product.

The perfume components useful herein may also be conveniently employed for deodorant purposes by incorporation into other products, for example, into laundry and household products such as rinse conditioners, household cleaners and detergent cleaners. The

perfume components can be incorporated into textiles themselves during their production using techniques known in the art, to provide deodorant protection.

In a preferred embodiment of the present invention, an Odour Reduction Value, measured in human axillae as described in Example 4, of at least 10%, more preferably at least 30%, and particularly at least 50% is obtained.

One or more of the perfume component(s) useful herein may be mixed with other perfume components e.g. perfume components of the prior art having deodorant properties, to formulate perfume compositions with desired deodorant and hedonistic properties.

In one such embodiment, there is provided a perfume composition as defined herein, wherein the perfume composition additionally comprises at least 15% by weight, preferably at least 30% by weight, of one or more of the following perfume components: acetyl di-iso-amylene, acetyl tributyl citrate, aldehyde C10 (i.e. decenal), Amber AB 358 (available from Quest International), amyl salicylate, anisyl acetate, Azarbre*, benzyl salicylate, cis-3-hexenyl salicylate, citral, citronellol, clove leaf distilled, coriander, cyclamen aldehyde, decen-1-ol, dihydroeugenol, diphenylmethane, Dupical*, Empetaal*, geraniol, helional i.e. 2-methyl-3-(3,4-methylene-dioxyphenyl)propanal, Ionones (alpha- and beta-), Jasmacyclene*, 3-(4-methyl-4-hydroxyamyl)-3-cyclohexene carboxaldehyde, methyl eugenol, methyl isoeugenol, Ortholate*, para-cresyl methyl ether, 2-phenylethyl alcohol, para tert. butyl cyclohexyl acetate, rose oxide (racemic), styrallyl acetate, tetrahydrolinalol, and vanillin; wherein all asterisked materials are trade marks of Quest International.

In a preferred embodiment, there is provided a perfume composition comprising:

- (i) at least 30% by weight of the perfume composition of at least 3 of the following perfume components: N-ethyl-N-(3-methylphenyl)propionamide, 2-ethyl-N-methyl-N-(3-methylphenyl)butanamide, dihydromyrcenol, (4-isopropylcyclohexyl)methanol, 3-methyl-5-phenylpentan-1-ol, 2,2,2-trichloro-1-phenylethyl acetate, isobornyl acetate, allyl amyl glycolate, alpha-terpineol, acetyl cedrene, tetrahydrogeraniol, citronellal, cuminic

aldehyde, cis-jasmone, pine American oil, peppermint (Chinese), 1,3,3-trimethyl-2-norbornanol, gamma-nonolactone, octahydro-2H-chromen-2-one, cis-4-decenal, 3-(3-isopropylphenyl)butanal; and

(ii) at least 30% by weight of the perfume composition of one or more of the following perfume components: acetyl di-iso-amylene, acetyl tributyl citrate, aldehyde C10, Amber AB 358, amyl salicylate, anisyl acetate, Azarbre, benzyl salicylate, cis-3-hexenyl salicylate, citral; citronellol, clove leaf distilled, coriander, cyclamen aldehyde, decen-1-ol, dihydroeugenol, diphenylmethane, Dupical, Empetaal, geraniol, helional, alpha-ionone, beta-ionone, Jasmacyclene, 3-(4-methyl-4-hydroxyamyl)-3-cyclohexene carboxaldehyde, methyl eugenol, methyl isoeugenol, Ortholate, para-cresyl methyl ether, 2-phenylethyl alcohol, para tert. butyl cyclohexyl acetate, rose oxide, styrallyl acetate, tetrahydrolinalol, and vanillin.

Also included within the scope of the invention is a method, particularly a cosmetic method, for reducing or preventing body malodour by topically applying to human skin a composition comprising a perfume component selected from at least one of the following; Armoise Tunisian oil, para-tert.butylphenylacetone nitrile, dihydrolinalol, N-ethyl-N-(3-methylphenyl)propionamide, 4-(5-ethylbicyclo[2.2.1]heptyl-2)-cyclohexanol, ethyltricyclo[5.2.1.0{2,6}]decane-2-carboxylate, 2-ethyl-N-methyl-N-(3-methylphenyl)butanamide, dihydromyrcenol, (4-isopropylcyclohexyl)methanol, 3-methyl-5-phenylpentan-1-ol, 2,2,2-trichloro-1-phenylethyl acetate, isobornyl acetate, allyl amyl glycolate, alpha-terpineol, acetyl cedrene, tetrahydrogeraniol, citronellal, cuminic aldehyde, 1,3,3-trimethyl-2-norbornanyl acetate, cis-jasmone, methyl octyl acetaldehyde, gamma-octalactone, octyl acetate, pine American oil, peppermint (Chinese), 1,3,3-trimethyl-2-norbornanol, gamma-nonolactone, octahydro-2H-chromen-2-one, cis-4-decenal, 3-(3-isopropylphenyl)butanal.

Preferably, the composition is a perfume composition.

Preferred perfume components for use in the method as defined above are selected from one or more of the following; Armoise Tunisian oil, para-tert.butylphenylacetone nitrile,

dihydrolinalol, N-ethyl-N-(3-methylphenyl)propionamide, 4-(5-ethylbicyclo[2.2.1]hept-2-yl)-cyclohexanol, ethyltricyclo[5.2.1.0{2,6}]decane-2-carboxylate, 2-ethyl-N-methyl-N-(3-methylphenyl)butanamide, (4-isopropylcyclohexyl)methanol, 3-methyl-5-phenylpentan-1-ol, 2,2,2-trichloro-1-phenylethyl acetate, isobornyl acetate, allyl amyl glycolate, acetyl cedrene, tetrahydrogeraniol, citronellal, cuminic aldehyde, 1,3,3-trimethyl-2-norbornanyl acetate, cis-jasmone, methyl octyl acetaldehyde, gamma-octalactone, octyl acetate, pine American oil, peppermint (Chinese), 1,3,3-trimethyl-2-norbornanol, gamma-nonolactone, octahydro-2H-chromen-2-one, cis-4-decenal, 3-(3-isopropylphenyl)butanal.

The method thus comprises topically applying to human skin, one or more of the specified perfume components which is(are) capable of reducing or preventing body malodour by inhibiting the production of odoriferous steroids by micro-organisms present on the skin surface, wherein the perfume component is capable of inhibiting the biotransformation of androstadienols to androstenones. Typically, the specified perfume components inhibit the production of odoriferous steroids by Coryneform bacteria present on the skin surface, particularly *Corynebacterium* sp. The inhibitory effect of the perfume components useful herein can be achieved antimicrobially or sub-lethally.

The antimicrobial effects of compounds, e.g. perfume components, are usually divided into two types; they can either inhibit bacterial growth (bacteriostatic action) or alternatively they can act by directly killing existing viable bacteria (bactericidal action).

The bacteriostatic action of a compound "X" such as a perfume component, can be tested for *in vitro* by inoculating a small number of bacteria into broths containing an appropriate range of concentrations of X. The broths are then incubated for a suitable time, and growth compared with a control containing no inhibitor. The broth containing the lowest concentration of X which shows reduction of growth compared to the control broth, is defined as the minimum inhibitory concentration (MIC).

The determination of bactericidal action of a compound "Y" such as a perfume component is carried out by adding various concentrations of compound Y to replicate broths

containing relatively high numbers of bacteria. After a certain period allowing any antibacterial activity to take place, aliquots of the bacterial cultures are diluted (usually in 10-fold steps) and dispensed onto agar plates. The plates are incubated with the expectation that each viable cell should produce a visible colony. The numbers of colonies are multiplied to take account of the dilution, to establish the number of viable cells in the broths. Once again, the broths containing compound Y are compared with an untreated control broth. The minimum concentration of compound Y which causes a reduction in the viable number of bacteria is the minimum bactericidal concentration (MBC). MBC can also be expressed in terms of the MBC required to produce a certain degree of killing (for example, a 3 log₁₀ reduction in count, equivalent to a 99.9% kill). Still further, the MBC can be expressed in kinetic terms - the time of exposure to an agent required for a given MBC effect.

A further possibility is that the process of inhibition could be sub-lethal (or sub-MIC), whereby the perfume components interfere with the metabolic process, but typically do not inhibit bacterial growth.

Preferably, the bacterial production of odoriferous steroids is reduced by at least 50%, more preferably by at least 70%, particularly by at least 80%, and especially by at least 90%. Three modes of achieving this reduction of odoriferous steroid production are possible. In the first mode, the perfume components (or perfume compositions) may act by direct (overt antimicrobial) killing of skin bacteria, e.g. by more than 10-fold; in the second mode, they may act on odoriferous steroid generation whilst maintaining a microbial cell viability of at least 70%; in the third mode, they may inhibit odoriferous steroid generation, at a concentration below the minimum inhibitory concentration (MIC), determined as described in Example 1 below. The third mode is preferred, since this provides malodour counteraction benefits, whilst leaving the natural skin microflora undisturbed. Thus, preferably the bacterial production of odoriferous steroids can be reduced or eliminated without significantly disturbing the skin's natural microflora. This may be achieved by inhibiting the bacterial enzymes responsible for the production of odoriferous steroids, in particular the androsthenones such as 5 α -androst-16-en-3-one.

In an even further aspect the present invention provides use of one or more of the following perfume components; Armoise Tunisian oil, para-tert.butylphenylacetonitrile, dihydrolinalol, N-ethyl-N-(3-methylphenyl)propionamide, 4-(5-ethylbicyclo[2.2.1]heptyl-2)-cyclohexanol, ethyltricyclo[5.2.1.0{2,6}]decane-2-carboxylate, 2-ethyl-N-methyl-N-(3-methylphenyl)butanamide, (4-isopropylcyclohexyl)methanol, 3-methyl-5-phenylpentan-1-ol, 2,2,2-trichloro-1-phenylethyl acetate, isobornyl acetate, allyl amyl glycolate, acetyl cedrene, tetrahydrogeraniol, citronellal, cuminic aldehyde, 1,3,3-trimethyl-2-norbornanyl acetate, cis-jasmone, methyl octyl acetaldehyde, gamma-octalactone, octyl acetate, pine American oil, peppermint (Chinese), 1,3,3-trimethyl-2-norbornanol, gamma-nonolactone, octahydro-2H-chromen-2-one, cis-4-decenal, 3-(3-isopropylphenyl)butanal, as a deodorant active.

Based on the MIC value evaluated for a particular perfume component, it is possible to select and combine those perfume components having low MIC values which are likely to be antimicrobially active, and to formulate a deodorant product which has some degree of anti-microbial activity. An example of this is a product including an antimicrobially effective amount, typically between 0.05% and 4% by weight, preferably between 0.1% and 2% by weight, more preferably between 0.5% and 1.5% by weight, of a perfume composition comprising at least 30% by weight of one or more of the following perfume components:

N-Ethyl-N-(3-methylphenyl)propionamide;
 3-Methyl-5-phenylpentan-1-ol;
 2,2,2-Trichloro-1-phenylethyl acetate;
 Pine American oil;
 cis-4-Decenal;
 4-(5-Ethylbicyclo[2.2.1]heptyl-2)-cyclohexanol;
 Cuminic aldehyde;
 Methyl octyl acetaldehyde;
 Armoise Tunisian oil;

Dihydromyrcenol;
 Allyl amyl glycolate;
 alpha-Terpineol;
 cis-Jasmone;
 Peppermint (Chinese);
 gamma-Nonalactone;
 Octahydro-2H-chromen-2-one;
 para-tert-Butylphenylacetonitrile;
 Dihydrolinalol;
 Tetrahydrogeraniol; and
 1,3,3-Trimethyl-2-norbornanol;

optionally in combination with perfume components having known high antimicrobial activity such as phenylethyl alcohol, geraniol, cinnamic acid, benzyl alcohol, and citral.

Likewise, it is also possible to select and combine those perfume components with higher MIC values which are likely to sub-lethally inhibit odoriferous steroid generation, and to formulate a deodorant product with minimal antimicrobial activity including, for example, appropriate levels of a perfume composition, typically between 0.05% and 4% of a perfume composition by weight of the deodorant product, preferably between 0.1% and 2% by weight, more preferably between 0.5% and 1.5% by weight, the perfume composition comprising at least 30% by weight of one or more of the following perfume components:

Ethyltricyclo[5.2.1.0{2,6}]decane-2-carboxylate;
 2-Ethyl-N-methyl-N-(3-methylphenyl)butanamide;
 (4-isopropylcyclohexyl)methanol;
 Isobornyl acetate;
 Acetyl cedrene;
 Citronellal;
 gamma-Octalactone;

Octyl acetate;

1,3,3-Trimethyl-2-norbornanyl acetate;

3-(3-isopropylphenyl)butanal.

The invention also provides the use of a perfume component to inhibit the biotransformation of androstadienols to androstenones, in particular the biotransformation of androsta-5,16-dien-3 β -ol to 5 α -androst-16-en-3-one.

The invention further provides the use of a perfume composition, comprising at least 30% by weight of one or more perfume components capable of inhibiting the biotransformation of androstadienols to androstenones, to reduce body malodour.

The invention further provides the use of a deodorant product, comprising a perfume component, to reduce body malodour by inhibiting the biotransformation of androstadienols to androstenones.

The invention still further provides a method of producing a perfume composition which comprises (i) evaluating perfume components on the ability to inhibit the biotransformation of androstadienols to androstenones, (ii) selecting perfume components on the ability to inhibit the biotransformation of androstadienols to androstenones, and (iii) mixing together two or more of said selected perfume components, optionally with other perfume components.

The invention still further provides use of a perfume composition comprising a perfume component to reduce body malodour, characterised in that the composition comprises at least 30% by weight of at least one of the perfume components specified in the paragraph bridging pages 5 and 6 above.

The invention is illustrated by the following examples.

Example 1: Minimum Inhibitory Concentration (MIC)

The minimum inhibitory concentration of a perfume component was determined by the following method.

A culture of the test strain - *Corynebacterium xerosis* NCTC 7243 (National Collection of Type Cultures, Public Health Laboratory Service, Central Public Health Laboratory, 61 Colindale Avenue, London) was grown in 100ml of tryptone soya broth (TSB) (Oxoid, Basingstoke, UK) for 16-24 hours, in a shaken flask at 37°C. The culture was then diluted in sterile 0.1% TSB (Oxoid, Basingstoke, UK) to give a concentration of bacteria of approximately 10^6 colony forming units (cfu) per ml.

Perfume or perfume component samples were diluted in sterile TSB. Each row of a standard, 96-well plastic microtitre plate (labelled A-H) was allocated to one sample, thus eight samples per plate. Row H contained only TSB for use as a bacterial control to indicate the degree of turbidity resulting from bacterial growth in the absence of any test material. Aseptically, 200µl of the initial dilution of perfume/perfume component was transferred to the 1st and 7th well of the appropriate row. All other test wells were filled with 100µl of sterile TSB using an 8-channel micro-pipette. The contents of each of the wells in column 1 were mixed by sucking samples up and down in pipette tips, before 100µl was transferred to column 2. The same sterile pipette tips were used to transfer 100µl of each well in column 7, into the appropriate well in column 8. This set of eight tips was then discarded into disinfectant solution. Using eight fresh, sterile tips the process was repeated by transferring 100µl from column 2 into column 3 (and 8 into 9). The process was continued until all wells in columns 6 and 12 contained 200µl. After mixing, 100µl was discarded from wells in columns 6 and 12 to waste. Finally, 100µl of pre-diluted bacterial culture (approx. 10^6 cfu/ml) was added, thus giving 200µl final volume in each well.

A blank plate was prepared for each set of eight samples in exactly the same way, except that 100 μ l of sterile 0.1% TSB was added instead of bacterial culture. Test and control plates were sealed using autoclave tape and incubated for 18-30 hours at 37°C.

The microtitre plate reader (Model MRX, Dynatech Laboratories) was preset to gently agitate the plates and mix the contents. The absorbance at 540nm (hereinafter referred to for brevity and simplicity as " A_{540} ") was used as a measure of turbidity resulting from bacterial growth. The control, un-inoculated plate for each set of samples was read first, and the plate reader then programmed to use the control readings to blank all other plate readings for the inoculated plates for the same set of test materials (i.e. removing turbidity due to perfume and possible colour changes during incubation). Thus, the corrected readings generated were absorbances resulting from turbidity from bacterial growth. The MIC was taken as the concentration of perfume/perfume component required to inhibit growth so that the change in absorbance during the incubation period was $< 0.2 A_{540}$.

Example 2: Steroid Biotransformation Assay

The ability of perfumery components and mixtures of these components to inhibit the biotransformation of androstadienols to androstenones was determined *in vitro* using the method described below.

Corynebacterium sp. NCIMB 41018 (National Collections Of Industrial, Food and Marine Bacteria, 23 St Machar Drive, Aberdeen, AB24 3RY, Scotland, UK) (also known as *Corynebacterium* G41) was grown in 100ml of TSB supplemented with 0.1% w/v yeast extract (Oxoid), and 0.1% v/v Tween 80 (Sigma, Poole, UK) for 18-30 hours, in a shaken flask at 37°C. This culture was then harvested by centrifugation, and resuspended in 100 ml of biotransformation medium (consisting of a sterile semi-synthetic basal medium containing KH_2PO_4 1.6 g/l; $(\text{NH}_4)_2\text{HPO}_4$ 5 g/l; Na_2SO_4 0.38 g/l; yeast nitrogen base 3.35 g/l; yeast extract 0.5 g/l; Tween 80 0.2 g/l; Triton X-100 0.2 g/l and $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ 0.5 g/l).


Substrate androsta-5,16-dien-3 β -ol (50mg/assay) was added to the bacterial suspension and incubated for 72 hours at 37°C with agitation (at 220-250rpm) in a 250 ml, baffled-Erlenmeyer flask.

Following biotransformation of androsta-5,16-dien-3 β -ol to androst-16-en-3-one the bacteria were harvested and the cell pellet dried in air and then under vacuum.

The dried cells were then crushed and suspended in a mixture of diethyl ether, chloroform, ethanol, ethyl acetate and acetone, and stirred for 16 hours. The supernatant was then reduced to half its volume, filtered and evaporated at 30°C and 15 mmHg pressure. The resulting residue was re-dissolved in 5 ml AR grade methanol. Following sonication, the sample was analysed by HPLC on a Phenomenex Luna 5 micron, C18 reverse-phase HPLC column coupled to a Millipore-Waters 600E System Controller. Elute was passed through a Millipore-Waters 486 Tuneable absorbance detector and relative amounts of the steroid metabolite was determined by a Hewlett Packard HP 3396A Integrator printer. The composition of the HPLC mobile phase was aqueous methanol. The flow rate was 0.8 ml/min. Calibration curves were used to determine the molar quantities of pure steroid metabolites in biotransformed mixtures and hence the conversions.

Metabolites were analysed by HPLC-MS to determine their structure.

The biotransformation of androsta-5,16-dien-3 β -ol to 5 α -androst-16-en-3-one by *Corynebacterium* NCIMB 41018 is as shown below:



It will be appreciated that the Coryneform bacteria used in Examples 1 and 2 are not the same strains. This is because the nutrient Tween-80 required for growth by *Corynebacterium* NCIMB 41018 (Example 2) is not suitable for inclusion in the growth medium used for MIC testing. As described above, during MIC testing, measurements are taken of the turbidity resulting from bacterial growth. Tween-80 when dissolved in an aqueous growth medium turns the medium cloudy. Thus, the addition of Tween-80 to a growth medium to be used for MIC testing would interfere with the readings, making an accurate determination of the turbidity due to bacterial growth impossible. Thus, a similar axillary *Corynebacterium* strain (*C. xerosis* NCTC 7243) is used in the MIC test, which does not require this nutrient for growth. The susceptibility of *Corynebacterium xerosis* NCTC 7243 a variety of perfume components is likely to be very similar to that of *Corynebacterium* NCIMB 41018 as they are from the same genus.

Example 3

Perfume A: Composition % by weight.

| INGREDIENT | w/w % | |
|---------------------------------|----------|---|
| AGARBOIS (Q) | 15 | * |
| CINNAMIC ALCOHOL | 2 | |
| COUMARIN | 1 | |
| DIHYDROMYRCENOL | 8 | * |
| GERANIUM OIL | 2 | |
| HABANOLIDE (F) | 3 | |
| LILIAL (G) | 10 | |
| (4-ISOPROPYLCYCLOHEXYL)METHANOL | 2 | * |
| MEFROSOL (Q) | 5 | * |
| METHYL ANTHRANILATE | 1 | |
| METHYL CEDRYL KETONE | 4 | |
| METHYL DIHYDROJASMONATE (Q) | 10 | |
| PHENYL ETHYL ALCOHOL | 15 | |
| ROSACETONE | 5 | * |
| VANILLIN 5% IN DEP | 17 - | |
| total | 100.00 % | |

* Materials of the invention

Trademarks: 'Q' = Quest International; 'F' = Firmenich; 'G' = Givaudan

Perfume B: Composition % by weight.

| INGREDIENT | w/w % | |
|---------------------------|-------|---|
| ACETYL CEDRENE | 7.5 | * |
| AGARBOIS (Q) | 6 | * |
| ALDEHYDE MNA 10% DEP | 1 | |
| ALLYL AMYL GLYCOLATE (Q) | 2.2 | * |
| AMBER CORE (Q) | 0.5 | |
| ARMOISE TUNISIAN | 0.4 | * |
| BANGALOL (Q) | 0.5 | |
| BENZYL SALICYLATE (Q) | 8.5 | |
| BERGAMOT OIL | 7.5 | |
| BOURGEONAL (Q) | 0.5 | |
| CARVONE LAEVO (Q) 10% DEP | 1 | |
| CEDARWOOD VIRGINIAN OIL | 1.1 | |
| cis-3-HEXENYL SALICYLATE | 1.5 | |
| CISTULATE (Q) 10% DEP | 2 | |
| CORIANDER | 0.3 | |

| | | |
|---------------------------------------|---------|---|
| COUMARIN | 0.6 | |
| CYCLOHEXYLOXYACETIC ACID, ALLYL ESTER | 0.2 | |
| CYCLOPENTADECANOLIDE | 2.2 | |
| DIHYDROMYRCENOL (Q) | 13 | * |
| ETHYLENE BRASSYLATE | 1.5 | |
| GERANIUM OIL | 1.4 | |
| HELIONAL | 0.3 | |
| HEXYL CINNAMIC ALDEHYDE | 2.5 | |
| IONONE (Q) | 1.5 | |
| ISO AMBOIS (Q) | 7.5 | |
| ISO BORNYL ACETATE | 0.6 | * |
| ISOBORNYL CYCLOHEXANOL | 1.5 | |
| LAVANDIN OIL | 0.3 | |
| LILIAL (G) | 6.8 | |
| METHYL CHAVICOL | 1.2 | |
| METHYL DIHYDROJASMONATE SUPER (Q) | 6.4 | |
| MOSS OAKMOSS SYNTHETIC | 0.2 | |
| NUTMEG PURE | 0.2 | |
| PEPPERMINT CHINESE 10% DEP | 3.5 | * |
| PETITGRAIN PARAGUAY | 0.2 | |
| ROSE OXIDE RACEMIC 10% DEP | 0.5 | |
| STYRALLYL ACETATE | 0.4 | |
| TERPINEOL ALPHA | 2.5 | * |
| TETRAHYDROLINALOL | 4.5 | |
| total | 100.00% | |

* Materials of the invention

Perfume C: Composition % by weight.

| INGREDIENT | w/w% | |
|-------------------------|------|---|
| ACETYL CEDRENE (Q) | 7 | * |
| AGARBOIS (Q) | 15 | * |
| ALDEHYDE MNA 10% DEP | 2.5 | |
| BENZYL SALICYLATE (Q) | 6.4 | |
| cis-JASMONE | 1.2 | * |
| CITRONELLAL | 2.2 | * |
| COUMARIN | 1.3 | |
| CYCLOPENTADECANOLIDE | 6.6 | |
| DIHYDROMYRCENOL (Q) | 8.5 | * |
| ETHYLENE BRASSYLATE | 2.3 | |
| HEXYL CINNAMIC ALDEHYDE | 3.5 | |

| | | |
|-----------------------------------|---------|---|
| ISO AMBOIS (Q) | 7 | |
| ISO BORNYL ACETATE | 2.6 | * |
| LILIAL (G) | 5.4 | |
| MARENIL (Q) | 1.3 | * |
| MEFROSOL (Q) | 5.4 | * |
| METHYL DIHYDROJASMONATE SUPER (Q) | 7.6 | |
| PETITGRAIN PARAGUAY | 1.2 | |
| TERPINEOL ALPHA | 3 | * |
| TETRAHYDROGERANIOL | 10 | * |
| total | 100.00% | |

* Materials of the invention

Perfume D: Composition % by weight.

| INGREDIENT | w/w% | |
|--|------|---|
| 4-(5-ETHYLBICYCLO[2.2.1]HEPTYL-2)-CYCLOHEXANOL | 1.2 | * |
| ACETYL CEDRENE (Q) | 5.3 | * |
| ALDEHYDE C11 (UNDECYLENIC ALDEHYDE) 10% DEP | 1.4 | |
| ALDEHYDE MNA 10% DEP | 0.8 | |
| ALLYL AMYL GLYCOLATE (Q) | 1.3 | * |
| ARMOISE TUNISIAN | 0.2 | * |
| BANGALOL (Q) | 0.3 | |
| BENZYL SALICYLATE (Q) | 5.1 | |
| BERGAMOT OIL | 4.8 | |
| CEDARWOOD VIRGINIAN OIL | 1.1 | |
| CITRONELLAL | 2 | * |
| CITRONELLOL | 6.9 | |
| CYCLOPENTADECANOLIDE | 2.3 | |
| DIHYDROMYRCENOL (Q) | 15.8 | * |
| ETHYLENE BRASSYLATE | 8.8 | |
| FENCHYL ACETATE | 2.5 | * |
| HEXYL CINNAMIC ALDEHYDE | 5.1 | |
| IONONE (Q) | 3.5 | |
| ISOBORNYL CYCLOHEXANOL | 1.8 | |
| METHYL DIHYDROJASMONATE SUPER (Q) | 5.5 | |
| PARA TERT BUTYL CYCLOHEXYL ACETATE | 3.4 | |
| PARADISAMIDE (Q) | 2.8 | * |
| PEPPERMINT CHINESE 10% DEP | 4.3 | * |
| PHENYLETHYL ALCOHOL | 6 | |
| ROSE OXIDE RACEMIC 10% DEP | 2.1 | |
| ROSEACETONE | 3.7 | * |
| TETRAHYDROGERANIOL | 2 | * |

total

100.00%

 * Materials of the invention

Perfume E: Composition % by weight.

| INGREDIENT | w/w% | |
|--|---------|---|
| 4-(5-ETHYLBICYCLO[2.2.1]HEPTYL-2)-CYCLOHEXANOL | 2.3 | * |
| AGARBOIS (Q) | 4 | * |
| ALDEHYDE C11 (UNDECYLENIC ALDEHYDE) 10% DEP | 1.2 | |
| AMBER CORE (Q) | 4.3 | |
| CARVONE LAEVO (Q) 10% DEP | 3.8 | |
| CEDARWOOD VIRGINIAN OIL | 1.8 | |
| cis-JASMONE | 0.5 | * |
| CISTULATE (Q) 10% DEP | 0.9 | |
| CITRONELLOL | 3.6 | |
| CORIANDER | 0.2 | |
| COUMARIN | 0.9 | |
| DIHYDROMYRCENOL (Q) | 4.5 | * |
| ETHYLENE BRASSYLATE | 6.2 | |
| FENCHYL ACETATE | 3.6 | * |
| HEXYL CINNAMIC ALDEHYDE | 6.8 | |
| HEXYL SALICYLATE | 7.5 | |
| LILIAL (G) | 6.5 | |
| MARENIL (Q) | 2.6 | * |
| METHYL CHAVICOL | 0.4 | |
| METHYL DIHYDROJASMONATE SUPER (Q) | 3.5 | |
| METHYL OCTYL ACETALDEHYDE 10% DEP | 5.5 | * |
| MOSS OAKMOSS SYNTHETIC | 0.2 | |
| PEPPERMINT CHINESE 10% DEP | 3.4 | * |
| PETITGRAIN PARAGUAY | 2.1 | |
| PHENYLETHYL ALCOHOL | 7.1 | |
| TERPINEOL ALPHA | 6.4 | * |
| TETRAHYDROGERANIOL | 8.2 | * |
| TETRAHYDROLINALOL | 2 | |
| total | 100.00% | |

* Materials of the invention

Example 4: Product Base Examples

The following are typical formulations of deodorant products which comprise a perfume composition in accordance with the invention. These formulations are made by methods common in the art.

1. Deodorant Sticks

| Ingredient | Content (% by weight) | |
|----------------------|-----------------------|----------------|
| | Formulation 1A | Formulation 1B |
| Ethanol | | 8.0 |
| Sodium Stearate | 7.0 | 6.0 |
| Propylene glycol | 70.0 | 12.0 |
| Perfume | 1.5 | 2.0 |
| PPG-3 Myristyl ether | | 28.0 |
| PPG-10 Cetyl ether | | 10.0 |
| Cyclomethicone | | 34.0 |
| Water | 21.5 | |

2. Aerosols

| Ingredient | Content % by weight | |
|----------------------|---------------------|----------------|
| | Formulation 2A | Formulation 2B |
| Ethanol B | up to 100 | |
| Propylene glycol | as required | |
| Perfume | 2.0 | 1.2 |
| Chlorhydrol microdry | | 31.8 |
| Silicone Fluid DC344 | | up to 100 |

| | | |
|-----------------|------|-------|
| Bentone gel IPP | | 13.65 |
| Dimethyl ether | 20.0 | |
| Concentrate | | 22.0 |
| Water | 23.0 | |

| Ingredient | Content % by weight |
|------------|---------------------|
| | Formulation 2C |

| | |
|---|-----------|
| Ethanol (Denatured) | up to 100 |
| Perfume | 1.0 |
| DC345 Fluid ⁽ⁱ⁾ | 15.0 |
| Hydrocarbon Propellant, 30 psig ⁽ⁱⁱ⁾ | 60.0 |

- (i) DC345 fluid (INCI name – CYCLOPENTA-SILOXANE) is a volatile, low viscosity, silicone fluid. It is non-greasy providing a light, silky feel on the skin.
- (ii) The hydrocarbon propellant can be any deodorised blend of n-butane, n-propane and isobutane having a pressure of 30 pounds per square inch gauge or 2.109 kg/cm² gauge (308 kPa).

3. Roll ons

| Ingredient | Content % by weight | |
|----------------|---------------------|----------------|
| | Formulation 3A | Formulation 3B |
| Ethanol | to 100% | 60.0 |
| Klucel MF | | 0.65 |
| Cremphor RM410 | | 0.5 |
| Perfume | 0.5 | 1.0 |
| AZTC* | 20.0 | |

| | | |
|----------------|------|-------|
| Cyclomethicone | 68.0 | |
| Dimethicone | 5.0 | |
| Silica | 2.5 | |
| Water | | 37.85 |

* Aluminium zirconium tetrachlorohydro glycinate

Perfume compositions A to E embodying this invention (see Example 3 above) were prepared and tested for deodorant action in underarm products, particularly an aerosol product of Formulation 2C, using an Odour Reduction Value test generally as described in US 4,278,658.

The Odour Reduction Value test was carried out using a panel of 40 Caucasian male subjects. A standard quantity (approximately 1.75g) of an aerosol product containing one of the perfume compositions or an unperfumed control was applied to the axillae of the panel members in accordance with a statistical design.

After a period of five hours and again at twenty four hours, the underarm odour was judged by three trained female assessors who scored the odour intensity in accordance with a 0 to 5 scale, as shown below:

| Score | Odour level | Conc. of aqueous isovaleric acid (ml/l) |
|-------|-------------|---|
| 0 | No odour | 0 |
| 1 | Slight | 0.013 |
| 2 | Definite | 0.053 |
| 3 | Moderate | 0.22 |
| 4 | Strong | 0.87 |
| 5 | Very Strong | 3.57 |

Average scores for each test product and the control product were then determined. The

score for each test product was subtracted from the score for the control product and the reduction expressed as a percentage to give the Odour Reduction Value(%).

Perfume compositions 'A' to 'E' were all found to exhibit significant deodorant activity.

For example, Perfume 'A' contains 35% of perfume components of the invention. Excluding diluents, this percentage increases to 42.2%. %. For this perfume, present at 1.0% in an aerosol product of Formulation 2C above, the Odour Reduction Value(%) compared to an unperfumed control was 66.9% (5 hours) and 33.8% (24 hours).

C245.00/Q

CLAIMS

1. A perfume composition comprising at least 30% by weight of one or more of the following perfume components; Armoise Tunisian oil, para-tert.butylphenylacetonitrile, dihydrolinalol, N-ethyl-N-(3-methylphenyl)propionamide, 4-(5-ethylbicyclo[2.2.1]heptyl-2)-cyclohexanol, ethyltricyclo[5.2.1.0{2,6}]decane-2-carboxylate, 2-ethyl-N-methyl-N-(3-methylphenyl)butanamide, dihydromyrcenol, (4-isopropylcyclohexyl)methanol, 3-methyl-5-phenylpentan-1-ol, 2,2,2-trichloro-1-phenylethyl acetate, isobornyl acetate, allyl amyl glycolate, alpha-terpineol, acetyl cedrene, tetrahydrogeraniol, citronellal, cuminic aldehyde, 1,3,3-trimethyl-2-norbornanyl acetate, cis-jasmone, methyl octyl acetaldehyde, gamma-octalactone, octyl acetate, pine American oil, peppermint (Chinese), 1,3,3-trimethyl-2-norbornanol, gamma-nonolactone, octahydro-2H-chromen-2-one, cis-4-decenal, 3-(3-isopropylphenyl)butanal.

2. A perfume composition comprising at least 3 of the following perfume components; Armoise Tunisian oil, para-tert.butylphenylacetonitrile, dihydrolinalol, N-ethyl-N-(3-methylphenyl)propionamide, 4-(5-ethylbicyclo[2.2.1]heptyl-2)-cyclohexanol, ethyltricyclo[5.2.1.0{2,6}]decane-2-carboxylate, 2-ethyl-N-methyl-N-(3-methylphenyl)butanamide, dihydromyrcenol, (4-isopropylcyclohexyl)methanol, 3-methyl-5-phenylpentan-1-ol, 2,2,2-trichloro-1-phenylethyl acetate, isobornyl acetate, allyl amyl glycolate, alpha-terpineol, acetyl cedrene, tetrahydrogeraniol, citronellal, cuminic aldehyde, 1,3,3-trimethyl-2-norbornanyl acetate, cis-jasmone, methyl octyl acetaldehyde, gamma-octalactone, octyl acetate, pine American oil, peppermint (Chinese), 1,3,3-trimethyl-2-norbornanol, gamma-nonolactone, octahydro-2H-chromen-2-one, cis-4-decenal, 3-(3-isopropylphenyl)butanal.

3. A perfume composition according to claim 2, wherein the perfume composition comprises at least 30% by weight of at least 3 of the specified perfume components.

4. A deodorant product comprising a perfume composition according to any one of claims 1, 2 or 3.

5. Use of one or more of the following perfume components; Armoise Tunisian oil, para-tert.butylphenylacetonitrile, dihydrolinalol, N-ethyl-N-(3-methylphenyl)propionamide, 4-(5-ethylbicyclo[2.2.1]heptyl-2)-cyclohexanol, ethyltricyclo[5.2.1.0{2,6}]decane-2-carboxylate, 2-ethyl-N-methyl-N-(3-methylphenyl)butanamide, (4-isopropylcyclohexyl)methanol, 3-methyl-5-phenylpentan-1-ol, 2,2,2-trichloro-1-phenylethyl acetate, isobornyl acetate, allyl amyl glycolate, acetyl cedrene, tetrahydrogeraniol, citronellal, cuminic aldehyde, 1,3,3-trimethyl-2-norbornanyl acetate, cis-jasmone, methyl octyl acetaldehyde, gamma-octalactone, octyl acetate, pine American oil, peppermint (Chinese), 1,3,3-trimethyl-2-norbornanol, gamma-nonolactone, octahydro-2H-chromen-2-one, cis-4-decenal, 3-(3-isopropylphenyl)butanal, as a deodorant active.

6. Use of one or more of the following perfume components; Armoise Tunisian oil, para-tert.butylphenylacetonitrile, dihydrolinalol, N-ethyl-N-(3-methylphenyl)propionamide, 4-(5-ethylbicyclo[2.2.1]heptyl-2)-cyclohexanol, ethyltricyclo[5.2.1.0{2,6}]decane-2-carboxylate, 2-ethyl-N-methyl-N-(3-methylphenyl)butanamide, (4-isopropylcyclohexyl)methanol, 3-methyl-5-phenylpentan-1-ol, 2,2,2-trichloro-1-phenylethyl acetate, isobornyl acetate, allyl amyl glycolate, acetyl cedrene, tetrahydrogeraniol, citronellal, cuminic aldehyde, 1,3,3-trimethyl-2-norbornanyl acetate, cis-jasmone, methyl octyl acetaldehyde, gamma-octalactone, octyl acetate, pine American oil, peppermint (Chinese), 1,3,3-trimethyl-2-norbornanol, gamma-nonolactone, octahydro-2H-chromen-2-one, cis-4-decenal, 3-(3-isopropylphenyl)butanal, in the manufacture of a composition for reducing or preventing body malodour.

7. A method for reducing or preventing body malodour by topically applying to human skin a composition comprising a perfume component selected from at least one of the following; Armoise Tunisian oil, para-tert.butylphenylacetonitrile, dihydrolinalol, N-ethyl-N-(3-methylphenyl)propionamide, 4-(5-ethylbicyclo[2.2.1]heptyl-2)-cyclohexanol, ethyltricyclo[5.2.1.0{2,6}]decane-2-carboxylate, 2-ethyl-N-methyl-N-(3-

methylphenyl)butanamide, dihydromyrcenol, (4-isopropylcyclohexyl)methanol, 3-methyl-5-phenylpentan-1-ol, 2,2,2-trichloro-1-phenylethyl acetate, isobornyl acetate, allyl amyl glycolate, alpha-terpineol, acetyl cedrene, tetrahydrogeraniol, citronellal, cuminic aldehyde, 1,3,3-trimethyl-2-norbornanyl acetate, cis-jasmone, methyl octyl acetaldehyde, gamma-octalactone, octyl acetate, pine American oil, peppermint (Chinese), 1,3,3-trimethyl-2-norbornanol, gamma-nonolactone, octahydro-2H-chromen-2-one, cis-4-decenal, 3-(3-isopropylphenyl)butanal.

8. A method according to claim 7 wherein the perfume component is selected from at least one of the following; Armoise Tunisian oil, para-tert.butylphenylacetonitrile, dihydrolinalol, N-ethyl-N-(3 methylphenyl)propionamide, 4-(5-ethylbicyclo[2.2.1]heptyl-2)-cyclohexanol, ethyltricyclo[5.2.1.0{2,6}]decane-2-carboxylate, 2-ethyl-N-methyl-N-(3-methylphenyl)butanamide, (4-isopropylcyclohexyl)methanol, 3-methyl-5-phenylpentan-1-ol, 2,2,2-trichloro-1-phenylethyl acetate, isobornyl acetate, allyl amyl glycolate, acetyl cedrene, tetrahydrogeraniol, citronellal, cuminic aldehyde, 1,3,3-trimethyl-2-norbornanyl acetate, cis-jasmone, methyl octyl acetaldehyde, gamma-octalactone, octyl acetate, pine American oil, peppermint (Chinese), 1,3,3-trimethyl-2-norbornanol, gamma-nonolactone, octahydro-2H-chromen-2-one, cis-4-decenal, 3-(3-isopropylphenyl)butanal.

9. A method according to claim 7 or 8, wherein the composition is a perfume composition.

10. A method according to claim 9, wherein the perfume composition comprises at least 30% by weight of at least one of the specified perfume components.

C245.00/Q

Title: Improvements in or relating to perfume compositions

Abstract

A perfume composition comprising a perfume component capable of inhibiting the production of odoriferous steroids by micro-organisms on the skin. The perfume component is capable of inhibiting the biotransformation of androstadienols to androstenones.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.